



# The Effectiveness of Mass Communication to Change Public Behavior

Lorien C. Abroms<sup>1</sup> and Edward W. Maibach<sup>2</sup>

<sup>1</sup>Department of Prevention and Community Health, George Washington University School of Public Health & Health Services, Washington, DC, 20037; email: [lorien@gwu.edu](mailto:lorien@gwu.edu)

<sup>2</sup>Center of Excellence in Climate Change Communication Research, George Mason University, Fairfax, Virginia 22030; email: [emaibach@gmu.edu](mailto:emaibach@gmu.edu)

Annu. Rev. Public Health 2008. 29:16.1–16.16

The *Annual Review of Public Health* is online at <http://publhealth.annualreviews.org>

This article's doi:  
10.1146/annurev.publhealth.29.020907.090824

Copyright © 2008 by Annual Reviews.  
All rights reserved

0163-7525/08/0421-0001\$20.00

## Key Words

mass media, ecological models, people and places framework

## Abstract

This article provides an overview of the ways in which mass communication has been used—or can be used—to promote beneficial changes in behavior among members of populations. We use an ecological perspective to examine the ways in which mass media interventions can be used to influence public behavior both directly and indirectly. Mass media interventions that seek to influence people directly—by directly targeting the people burdened by the public health problem of concern and/or the people who influence them—have a long basis in public health history, and recent reviews have clarified our expectations about what can be expected from such approaches. Mass media interventions that seek to influence people indirectly—by creating beneficial changes in the places (or environments) in which people live and work—have equal if not greater potential to promote beneficial changes in population health behaviors, but these are currently less explored options. To have the greatest possible beneficial influence on public behavior with the public health resources available, we recommend that public health program planners assess their opportunities to use media to target both people and places in a manner that complements and extends other investments being made in population health enhancement.

## INTRODUCTION

Over the past two decades—since our first contribution to the *Annual Review of Public Health* (24)—the world has evolved in dramatic ways with regard to mass communication and public health. A communication revolution occurred that has blurred the traditional distinctions between mass and interpersonal communication, and it changed in many ways how we must think about using the media to promote the public's health. This revolution was driven by two primary factors. The first of these factors was the extraordinary proliferation of media channels and the consequent decline in media audience sizes (43). In the late 1980s, mass media were thought of as “broadcast” media. Now, mass media channels are more aptly described as “narrowcast” or even “slivercast” media because of significantly reduced audience sizes and because of the financial need for each channel to focus narrowly on the viewing or reading interests of a specific demographic or psychographic audience (42). It is therefore more difficult for public health organizations to reach “mass” audiences through the media because doing so now requires using limited public health communication resources across a wide array of channels.

A second major factor driving the communication revolution is the rise of the Internet (77, 78). The Internet and the World Wide Web are quintessential disruptive technologies in that they radically altered—and continue to alter—both the ways in which information flows through our society and the business models that support the mass media industry (11, 77). Until the mid 1990s, interpersonal communication occurred largely on a one-to-one or a one-to-few basis; the growth of the Internet and its myriad manifestations (including blast emails, Web sites, blogs, and RSS feeds) has given individuals and organizations dynamic and powerful new communication tools by which to spread their influence directly to thousands—indeed millions—of people independent of traditional mass me-

dia channels (and their associated gatekeepers). The ascendancy of nonbroadcast communication platforms is an extremely positive development for the field of public health because it offers us many powerful and flexible communication tools through which to convey important information to a variety of audiences. Moreover, these new communication tools can be inexpensive compared with the costs of traditional mass media outreach (77).

The past two decades have also witnessed a dramatic revolution in how we in public health understand population health. The rise of ecological models of health has helped us appreciate more clearly that the determinants of population health exist on multiple levels and include not only the characteristics of individuals (e.g., their attitudes and beliefs), but also the characteristics of social networks, the organizations in people's lives (e.g., workplace, church, sports league), and the neighborhoods in which they live (4, 41). Ecological models of health make clear that to use public health resources efficiently, we must make efforts across multiple levels of influence to affect the full range of factors that undermine—or promote—population health (44, 48, 66, 76).

The aftermath of a revolution—or in this case, the aftermath of two concurrent revolutions—is an excellent time to revisit one's operating assumptions. We recently did just that and concluded that the communication and the public health revolutions have not yet been adequately integrated into a new understanding of, or framework for, public health communication. In an article titled “Communication and Marketing as Tools to Cultivate the Public's Health: A Proposed ‘People and Places’ Framework,” we suggested a new way of understanding and pursuing the potential of communication interventions to change public behavior and promote the public's health (44).

This purpose of this review is to explore the implications of an ecological perspective on mass media campaigns as a means to change

public behavior. Using our recently developed ecological framework, the people and places framework (44), we review the degree to which and the ways in which mass media campaigns have targeted various levels of influence or fields of influence of the ecological framework. Where possible, we identify factors associated with successful campaigns within each field of influence. We make the case that the vast majority of public health media campaigns to date have targeted only a small range of ecological factors, thereby limiting our understanding of the true potential for mass media interventions.

## BACKGROUND

For a while, various experts who have reviewed the health communication campaigns literature, or some aspect of that literature, have tended to reach similar conclusions. Namely, most have concluded that mass media interventions, by themselves or in combination with other programs, can significantly influence the health behaviors of populations (5, 24, 39, 52, 59, 63, 72, 59, 92). One important caveat is associated with this conclusion, however: The effects of health communication campaigns are typically only modest in size (52, 72). Although there are clear exceptions to this rule—i.e., campaigns that have had dramatic behavior change impacts (21, 33, 49, 71) as well as campaigns that have had no behavior change impact (25, 64)—the rule itself apply to a broad range of public health media campaigns.

Effective public health media campaigns typically have two important qualities: They feature well-designed messages, and those messages are delivered to their intended audience with sufficient reach and frequency to be seen or heard and remembered (33, 52, 59). While the art and science of effective health message design continues to develop (70), the importance of following these principles has been well understood—and has been the subject of considerable attention in the field of public health—for quite some time (45, 63).

Indeed, Hornik (2002) noted that the public health communication field has been perhaps too focused on issues of message design and not adequately focused on the more costly challenge associated with achieving sufficient levels of message exposure (i.e., reach and frequency) among members of the target audience (33).

Extant reviews of the public health communication literature are limited in an important way. Most of what we know about the potential of public health media campaigns comes from campaigns that sought to influence population behavior by targeting individual-level antecedents to the behavior of concern (such as knowledge, perceptions, and self-efficacy). As such, extant literature reviews can reveal only a constrained view of the potential of public health communication. A more complete view—one that is in better sync with contemporary thinking in public health—requires that we gain an understanding mass media campaign potential across the full range of factors implied by ecological models of health.

Ecological models of health consider both the characteristics of individuals and the contexts in which they live (76). Following from this perspective, we recently proposed the people and places framework as a simple ecological model that posits that the populations' health is influenced by (a) the attributes of the people in the population, (b) the attributes of the environments or places in which members of the population live, work, go to school, shop,; and (c) important interactions between the attributes of people and places (44). These attributes and their interactions influence health through their impact on health behavior and their effects on physical functioning and well-being (4). The framework is illustrated in **Figure 1** (see color insert).

In this framework, the attributes of people can be understood through three levels of analysis, or as we have termed them, three fields of influence: the individual field; the social network field; and the group, community,

or population field. In addition, building on the important work of Farley & Cohen (19), the attributes of places can be understood in terms of two fields of influence: local and distal fields of influence. Local fields of influence are those in a person's immediate environment: within his/her own home, neighborhood, and workplace. Distal fields of influence are those further afield, but that still make an impact on a person's life. For example, decisions made in distal places such as the national capital or by multinational corporations exert influence over the behavior and health of people who live over wide geographic areas.

### USING MASS MEDIA INTERVENTIONS TO CULTIVATE HEALTH IN EACH FIELD OF INFLUENCE

Mass media interventions can be, and to a lesser extent have been, used to influence functioning in each of the five fields of influence identified in our people and places framework. To add a new perspective to the results of published literature reviews and meta-analyses, as well as to inform our own points of view, we searched the peer-reviewed literature to assess the prevalence and find examples of reported mass media campaigns targeting variables in each of the five fields of influence during the most recent decade (1997–2007). For the purposes of this review, we defined mass media campaigns as any planned effort that disseminates messages to produce awareness or behavior change among an intended population through channels that reach a broad audience (5). These channels can include traditional media such as radio, television, magazines, billboards, and newspapers, as well as those that make use of newer technologies such as email, cell phones, and interactive Web sites.

#### Influencing Individual-Level Factors

The effects of public health communication interventions on individual-level factors that

influence health behaviors have been the subject of research for many decades. The literature points to the following individual-level factors as being predictive of health behaviors: cognitions (e.g., knowledge and beliefs, self-efficacy, and outcome expectancies) (1), affect (e.g., depression) (38), skills (e.g., contraceptive skills) (8), motivation (e.g., high intrinsic interest) (53), and intentions (23). Mass media campaigns can and have sought to influence these factors for their own sake, and as a means to change behavior. Biological predispositions (e.g., sensation seeking) (55) and demographic factors (e.g., gender, income, employment status) (46) are additional individual-level factors that can and have served as a means to stratify (i.e., segment) audiences and target messages.

The vast majority of campaigns that we identified in our literature review targeted individual-level factors. Some of these campaigns have been spectacularly successful. One of the most successful examples of a public health communication campaign targeting individual-level factors is the “truth” campaign as conducted first by the Florida Department of Health, and then by the American Legacy Foundation. This campaign is aimed at youth between the ages of 12 and 17 years. Its ads—which often feature trendy youth involved in public demonstrations against the tobacco industry—are intended to provide tobacco-prone adolescents with the “truth” about the deceitful marketing practices of the tobacco industry. The ads encourage youth to rebel against the tobacco industry instead of rebelling with tobacco (31). The national campaign has been consistently associated with an increase in antitobacco attitudes and beliefs (21) and is responsible for 22% of the observed decline in youth smoking between 1999 and 2002 (20). The earlier campaign in Florida was even more successful, presumably because of a significantly larger per capita budget (71). Similar-themed campaigns have also been successfully employed in California and Massachusetts (56, 69).

As noted above, existing reviews of mass media campaigns have focused largely on

campaigns that target individual-level factors. From these campaigns we know that large campaign effects on individual-level factors appear to be the exception rather than the rule. Modest to moderate effects—when the campaigns are well designed and of adequate intensity—are the rule (52). Snyder & Hamilton's (72) meta-analysis of 48 published community-wide mass media health campaigns—all of which were conducted in the United States and were evaluated with a quasi-experimental research design—successfully quantifies the size of typical campaign effects. Overall, the effect size of the average campaign on population behavior in the short-term, as measured by the mean of correlations, was 0.09, which “roughly translates to 9% more people performing the behavior after the campaign than before” (72). The average effect size for campaigns that promoted behaviors that were enforceable by law (e.g., seat belt use) was considerably higher (0.17), whereas the effect size for purely “persuasive campaigns”—defined as those campaigns not promoting a legally enforceable behavior—was considerably smaller (0.05). This meta-analysis also identified two important variables that moderated the behavioral influence of media campaigns: reach and novelty of the information presented. Campaigns that achieved a higher reach (i.e., the proportion of the target audience who were exposed to the campaign) had a larger impact, as did campaigns that presented new information (vs. information that had already been communicated previously in other ways).

It goes without saying that nearly all contemporary mass media campaigns include the creation and promotion of a Web site as part of their collateral materials as a means to gain additional exposure to their messages. More sophisticated Web sites may include features such as interactive games or “advergames” (which are most common in campaigns targeting kids), downloads (e.g., campaign materials, screensavers, buddy icons), cell phone applications, and expert systems (i.e., an automated counselor that collects information

on the user and then guides him/her using tailored advice to the recommended behavior change).

Thus far, few published studies in the public health literature examine the contribution of a Web site or Web site components to campaign effectiveness. However, existing indicators are promising. First, the commercial sector's evidence documents the positive impact of Web-based promotions on customer reach, brand awareness, customer relationship building, and product sales (10). Second, investigators are seeing a growing number of cases of Web-based technologies being successfully used to generate high levels of campaign reach and engagement. For example, CDC's VERB campaign, a national campaign aimed at increasing physical activity among children aged 9–13 years, which began in 2002, generated more than 10 million independent visits to its Web site, <http://www.VERBnow.com>, over the course of 2 years. Of these visits, more than 1.1 million 9–13 year olds—or 5.3% of all American tweens (83)—registered with the Web site and recorded their hours of physical activity or searched for places in their zip code to be active (M.E. Huhman, personal correspondence, 2007). Although the unique contribution of these efforts to the high level of overall campaign awareness and to the observed increases in physical activity remain unclear (35), the total number of people reached and engaged is impressive. Third, an emerging body of evidence in the public health literature indicates that Web-based expert systems—independent of supporting media campaigns—can have impressive effects on individual health beliefs, attitudes, and behavior (77, 78).

### Social Network Level

An individual's web of social ties can be defined as his/her social network (22). Compelling evidence demonstrates that being embedded in a large social network positively affects health. Studies have consistently shown that individuals who lack social ties are

less likely to adopt recommended health behaviors and are less healthy overall, psychologically and physically (3, 4, 34). Various aspects of social networks have been identified that can be health promoting. These include, at a minimum, size and connectedness of a person's social network, diversity of ties in the social network, the degree to which the various relations in a social network (e.g., parents, friends, teachers, and mentors) provide social support, positive modeling, guidance and monitoring (4, 34, 82), and the presence of positive health opinion leaders in the social network (17, 62, 84).

The literature shows many examples of mass media campaigns that have been used to stimulate beneficial changes in social network-level factors. The preponderance of these media campaigns have targeted members of the inner circle of an individual's social network as a means to influence the ultimate target audience—those people who are most affected by the public health problem of concern. Most of these campaigns are aimed at good friends (2, 50), parents (16, 54, 79), older siblings (54), and spouses (68). These campaigns, for the most part, ask social network members to provide guidance and monitoring (54) and to encourage the other person to adopt various health behaviors [e.g., abstain from unprotected sex (16, 79), do not smoke (2), and do not drive drunk (50)]. To a lesser extent, these campaigns have attempted to increase the provision of social support among network members (68).

One example of a successful campaign, which targeted parents as a means to influence adolescents, was the North Carolina Department of Public Health's effort to prevent teenage pregnancy. With the tagline, "Talk to your kids about sex. Everyone else is," parents were urged through TV, radio, and billboard ads to talk to their children about safe sexual behavior. On the basis of a phone survey of parents, parents exposed to the campaign were more likely to have recently talked to their adolescent children about sex and more likely to talk to their children in the next month about sex (16).

The use of mass media to stimulate interpersonal communication—that is, to encourage members of social networks who are exposed to a campaign to discuss the topic of the campaign with others in their social network and thereby pass on or reinforce the prescribed health information or practice—is another strategy that has been productively used. Some evidence suggests that campaigns that can stimulate interpersonal communication about the campaign topic generate larger behavior change effects than do campaigns lacking this effect (7, 29, 85). This was true in a family-planning media campaign in Nepal. Women exposed to campaign messages through conversations with others were more likely to adopt the recommended behavior of contraceptive use than were those who were only directly exposed to the mass media campaign (7).

Although the large majority of media campaigns targeting social networks have focused on influencing existing social ties, a limited number of examples of mass media campaigns set out to increase the size of social networks. One example of such a campaign is the Harvard mentoring project (see sidebar, Listing of Web Sites of Ongoing Campaigns). This mass media campaign was designed to increase the social networks of underprivileged children by encouraging adults to volunteer to be mentors to these children. The campaign used a three-pronged strategy: public service announcements, outreach to the

### LISTING OF WEB SITES OF ONGOING CAMPAIGNS

Harvard Mentoring Project:

<http://www.hsph.harvard.edu/chc/mentoring.html>

Live Earth concert series: <http://www.liveearth.org/>

Rock the Vote: <http://www.rockthevote.com/>

Smoke Free Movies: <http://smokefreemovies.ucsf.edu/>

The truth campaign: <http://www.thetruth.com/>

The VERB campaign: <http://www.VERBnow.com>

entertainment community (i.e., efforts to get appropriate storylines developed in entertainment media), and news media outreach. This campaign, which is ongoing, has been successful in gaining significant donated airtime and in generating more than 700,000 calls to a hotline from people interested in mentoring (9).

To our knowledge, the impact of social network-level public health communication campaigns, in terms of their success in influencing social network-level variables or public behavior, has not been systematically reviewed. Most social network-level mass media interventions have focused on attempting to harness the influence of a person's inner coterie of friends and family. We need to explore the other aspects of social networks, those that are health promoting and potentially amenable to change through the mass media.

### Community Level

Characteristics of communities that might be targets for mass media campaigns include social norms, social capital, social cohesion, collective efficacy, income inequality and racism. Social cohesion can be defined as the extent of connectedness and solidarity within a group (4). Highly cohesive communities are thought to be endowed with large stocks of social capital, or characteristics that facilitate collective action such as interpersonal trust, norms of reciprocity, and mutual aid (57). Social capital has been linked to self-reported health and mortality (40, 41), as well as to the recollection of health information (87). A similar relationship has been established between a community's level of income inequality—the income gap between the most well off and least well off—and health, whereby communities with higher levels of income inequality are associated with higher levels of morbidity and mortality (58, 93).

Without question, the most common use of media campaigns to influence community-level factors is aimed at altering perceived

social norms. The literature richly documents that social norms play a powerful role in shaping the behaviors of people in populations (18); many people find it difficult to behave in a manner inconsistent with the social norm, even if the norm is not something to which they particularly subscribe. Social norms media campaigns are based on the observation that certain risk behaviors, particularly among adolescents and young adults, are based in part on a mistaken perception of the social norm (14). Many of these social norms campaigns have taken place in college settings and targeted norms pertaining to alcohol consumption because college students greatly overestimate the levels of heavy drinking among their peers (6). In these cases, the mass media campaign is used as a tool to correct misperceptions about normative levels of college drinking, and thereby attempt to reduce college alcohol consumption. Several studies evaluating social norms media campaigns demonstrate their efficacy (14, 27), although not all such campaigns have succeeded (65). The most rigorously designed of the studies found that, among 18 college campuses included in the study, those randomized to receive a social norms alcohol reduction campaign experienced greater reductions both in perceptions about normative levels of alcohol consumption and in actual drinking behavior (14).

The mass media has been used to increase social capital, social cohesion, or collective efficacy in only a handful of instances (28, 47, 80, 88). In one example, the Kansas Health Foundation conducted a statewide media campaign to increase social capital in Kansas (80). Over a two-year period, the Foundation released a series of paid television and print ads aimed at increasing social capital by encouraging nonparental adults to involve themselves in children's lives and in their community. As a result of the campaign, improvements were observed in attitudinal measures such as attending to youth and forming attachment to the community but not in behavioral outcomes such as having engaged with children

or organized other adults to address the needs of children (80). Another interesting example—although not directly from the field of public health—occurred during the 2004 U.S. presidential primary campaign. Rock the Vote, a national youth vote organization, partnered with CNN to sponsor a nationally televised debate where Democratic presidential candidates responded to questions posed directly by young citizens. Young viewers of this event experienced greater identification with the candidates and enjoyed a heightened level of civic engagement—a construct closely related to social capital—as compared with young viewers of a traditional journalist-led debate format (47). A third example occurred as we were finishing this article. The “Live Earth” concert series—a 24-hour, 7-continent concert televised series held on July 7, 2007—featured extensive messaging intended to promote an enhanced sense of collective efficacy around the issue of global warming. A prepost evaluation of this media event—reported to have been the largest televised music event ever—is currently being conducted (A. Leiserowitz, personal correspondence, 2007).

Clearly more work is needed to explore the potential of using mass media campaigns to cultivate health at the community level of influence. Many aspects of community—such as social capital, social cohesion, collective efficacy, income inequality, and racism—are important determinants of health that may be influenced in a cost-effective manner through mass media interventions.

### Place: Local and Distal Levels

Places—homes, schools, work sites, roads, grocery stores, neighborhoods, and cities—affect our health behaviors and health in a variety of complex and subtle ways. Thus it is often difficult to conceptualize the range of opportunities for intervening with the places that surround us. Cohen, Scribner and Farley (12) provided a useful system for categorizing the place-based field of influence into

four distinct subdomains. These consist of influences related to (a) the laws and policies in the environment, (b) the availability of products and services in the environment (such as health services, condoms, or fresh fruits and vegetables), (c) the physical structures in the environment (such as the presence or absence of sidewalks), and (d) the media and cultural messages in the environment (such as an abundance of unhealthy foods advertised on TV). Each of these distinct aspects of place represents potential targets for place-based mass media campaigns.

The mass media strategy most commonly used in changing the place-based field of influence is media advocacy. Media advocacy has been defined as “the strategic use of mass media in combination with community organizing to advance healthy public policies” (88). Media advocacy generally involves framing public health issues to emphasize the policy or environmental solution, gaining access to the news or other forms of the mass media as a means to reach the public and policy makers, and using this access to mobilize the public and force policy makers to enact particular policy solutions.

Only a small number of well designed evaluations of media advocacy efforts have been conducted, but those studies indicate that media advocacy efforts are promising (30, 32, 51, 75). One such study evaluated the media advocacy activities of the Florida Tobacco Control Program. The program aims to garner media coverage to encourage counties across Florida to adopt product placement ordinances for tobacco products (e.g., tobacco products must be kept behind the counter of a store). As a result of the campaign, newspaper stories about the program’s policy activities increased relative to other tobacco control topics covered during the intervention period. Furthermore, counties that experienced more newspaper coverage of the program’s activities were more likely to adopt the targeted new tobacco product placement ordinances. Unfortunately, the new ordinances, once they went into effect, did not result in the expected

declines in youth smoking (51). Some positive effects have been observed for media advocacy efforts in the arena of clean indoor air legislation (60) and for policies related to reducing alcohol abuse and alcohol-associated fatalities (32), although, for alcohol, effects have been mixed (30).

Although most published studies document media advocacy efforts to change governmental laws and policies (which often address the availability of products and services), mass media campaigns have also been used to directly alter the content of the media messages present in our environment. Compelling evidence shows that our media environment—both the advertising and the content of programming—shapes our health behaviors (86). One notable example of using the mass media to alter the media environment is an effort led by Smokefree Movies. Smokefree Movies is a group that strives to ensure that children will not be exposed to depictions of smoking in movies, an established risk factor for smoking uptake (67). Through a series of advertisements published in *Variety*, a movie industry publication, as well as in high-profile newspapers such as the *New York Times*, Smokefree Movies has pressured the Motion Picture Association of America (MPAA) to give films with smoking scenes an R rating (26), in much the same way that R ratings are given to films with foul language or sexually explicit scenes. Although there is no formal evaluation of the six-year campaign, recent news events indicate that the group is making some headway. On May 10, 2007, the MPAA announced that it was revising its rating system to include smoking in movies as a factor when assigning ratings to films, a step in the right direction, although the action fell short of Smokefree Movies' stated goal of a mandatory R rating (81).

Media advocacy has been heralded over the past decade as an important public health strategy and has been used relatively frequently (15, 88, 89). Although examples similar to the campaign set out by Smokefree Movies suggest positive effects, the evidence

base to support media advocacy as an effective public health strategy is surprisingly thin (73, 74). Relatively few studies have been published that involve media advocacy, and of those published, most are descriptive and rely on a case history analysis to explain its impact (74). One major factor driving this paucity of published studies is the lack of established and well-developed methodologies for evaluating media advocacy and other such complex environmental interventions (73, 74). Tackling the methodological hurdles for media advocacy is an important first step needed to strengthen our understanding of the value of media advocacy for shaping the place-based field of influence.

## DISCUSSION

Many people and thought-leading organizations in the field of public health are fundamentally optimistic about our ability to harness the potential of mass media to promote the public's health. The Institute of Medicine, for example, has issued a number of recent reports that indicate their strong belief in the power of media and communication as a public health strategy (36, 37). Indeed, in our highly saturated media environment—in which the average American adult spends more than 10 h and the average American child spends more than 6 h per day consuming media—there are myriad opportunities to reach audiences with communication interventions (43, 61). The evidence is fairly compelling that interventions targeting individual-level factors can be a highly cost-effective way to promote population health. Skeptics of the prospects for public health communication intervention—and many well-grounded skeptical assessments have taken place over the past quarter century (88, 90, 91)—however, are likely to conclude that this exuberance is irrational given the small population effects of individual-level public health media campaigns.

In the most thorough assessment to date (i.e., an edited volume seeking to gain

perspective on the potential of public health communication campaigns), Hornik (2002) concluded that ample evidence support an optimistic view of public health communication, although he noted that the very nature of highly effective public health communication programs tends not to be easily evaluated with traditional methods (33). Hornik alluded to “big messy programs” in describing the public health communication programs that have had the biggest impact. This term, “big messy program,” describes communication interventions not as a precisely targeted (almost surgical) intervention aimed at individuals, but rather as a program that includes many diverse communication tactics (e.g., mass media messages, interpersonal communication, and outreach to policy makers) often directed at changing both individuals and the social system. In contrast, he concluded that the mass media campaigns most likely to fail (i.e., those that do not influence public behavior) are those aimed solely at changing individuals. In Hornik’s opinion, these programs fail precisely because they do not change the larger social system and are therefore unable to achieve adequate contact (i.e., message reach and frequency) with their intended audience.

We share Hornik’s and the Institute of Medicine’s optimistic view of public health media campaigns; their true potential can be better understood, and pursued, by embracing an ecological framework of health that considers determinants of health across multiple fields of influence. Informed by the people and places framework, individual-level campaigns may fail to deliver the big results because they, in a variety of ways, encourage and help individuals to change their behavior, but ignore the many social-network, community-level, and place-based barriers to change. Although public behavior does indeed change on average under these circumstances (72), it is not of sufficient magnitude to solve pressing public health problems. By harnessing the mass media to change social support, community norms, the availability of

products and services, and other factors from nonindividual fields of influence, we should be able to improve greatly the likelihood of achieving large-scale changes in public behavior.

We regrettably understand that this article is not a definitive review of the potential to use mass media to change public behavior. Because of the relatively few studies investigating the uses of mass media for changing social network-, community-, and place-based factors, the state of the literature is still too weak to write that review. Rather, we ask this question: How can we use communication to influence the full range of important factors suggested by ecological models of health? Some public health communication programs must be, by necessity, of a limited nature. Conversely, many can be of the “big messy program” variety and can target factors across multiple fields of influence. Regardless of which type of program is being designed, campaign planning will benefit from clearly assessing the factors across the fields of influence that are most influential in creating the status quo and by making informed decisions about how to use available communication research to target those factors.

At the most basic level, we advocate to move beyond the Either/Or mentality that has characterized the public health communication field for the past several decades. Mass media campaign resources are, without question, limited, but this does not mean that they should be invested either to change the health behaviors of people or to change the health-enhancing capacity of places. Both of these general approaches are worth pursuing, or at least worth considering. Some public health organizations, by virtue of their circumstances and resources, will decide to focus on one approach or the other, and this is a perfectly justifiable decision. The public health community as a whole, however, should not be investing in one approach or the other, but rather in using communication to cultivate change in both areas of influence.

The issue of micronutrient fortification of staple foods offers an illustration of our point. For the past few decades, public health professionals have sought to eliminate micronutrient deficiencies by encouraging or requiring the manufacturers of certain staple foods to fortify the food with the missing nutrient (e.g., vitamin A). Mass media campaigns have been used—or can be used—in a variety of ways to effect this change. Campaigns can be used to advocate to policy makers the benefits of requiring micronutrient fortification as a matter of law. Campaigns can be used to target manufacturers directly to encourage them to voluntarily modify their manufacturing practices. And finally, campaigns can be used to target members of the affected population—and/or the people who influence them—to encourage them to seek and purchase only fortified food staples. A recent review of the literature concluded that the most effective micronutrient fortification programs are those that take measures both to ensure the wide availability of fortified foods and to take measures to create consumer demand for such foods (13).

A similar picture emerges from campaigns targeting the use of seatbelts in cars. As noted earlier, Snyder & Hamilton (72) found in their meta-analysis that campaigns promoting seatbelt use that emphasized enforcement messages—that is, messages about the fines or other legal consequences of not using a seatbelt (e.g., Click It or Ticket)—were found to be more than three times more effective than campaigns, seatbelt campaigns included, that relied solely on persuasive approaches. Furthermore, they reported that seatbelt laws in the absence of a media campaign were less effective than the combination of the seatbelt law with the enforcement-based media campaign (72). Apparently, it was the combination of a place-based strategy (in this case, the enforcement policy) and a people-based strategy (the mass media campaign targeting individual drivers) that made the greatest impact on behavior change.

Our assessment of the published research on mass media campaigns finds that the preponderance of evaluated interventions have targeted the individual field of influence, and to a lesser extent, the social network field of influence. It is not entirely clear why social network, community, and place-based factors have been underemphasized as targets for mass media interventions, but we would be wise to turn our attention to exploring systematically how we can use public health communication interventions to cultivate change in these fields of influence. There is clearly a need to review, synthesize, and if possible meta-analyze the extant public health communication literature that has sought to create change at the social network and the community or population level, as well as at the local and distal level of place. The history of our field is characterized by a focus on the individual, but certainly there is now enough research to begin to improve and expand our understanding of the potential of public health communication by assessing what we know about its potential to influence public behavior positively through other fields of influence. These studies can, and likely should, come both from a renewed effort to systematically evaluate the public health communication efforts of public health practice organizations as well as from sponsored research conducted specifically to advance our understanding of how communication can be used to shape public behavior positively.

At a deeper level, our peers in public health communication should consider using all the fields of influence suggested in the people and places framework to structure their analyses of intervention opportunities. Furthermore, we recommend that they develop mass media campaigns in a manner that complements and extends other investments being made in population health enhancement. Such a strategy will lead to the use of public health resources in a manner that is efficient, effective, and ethically responsible.

## DISCLOSURE STATEMENT

The authors are not aware of any biases that might be perceived as affecting the objectivity of this review.

## LITERATURE CITED


1. Bandura A. 2004. Health promotion by social cognitive means. *Health Educ. Behav.* 31:143–64
2. Bauman KE, LaPrelle J, Brown JD, Koch GG, Padgett CA. 1991. The influence of three mass media campaigns on variables related to adolescent cigarette smoking: results of a field experiment. *Am. J. Public Health* 81(5):597–604
3. Berkman L. 1985. The relationship of social networks and social support to morbidity and mortality. In *Social Support and Health*, ed. S Cohen, SL Syme, pp. 241–62. New York: Academic
4. Berkman LF, Glass T. 2000. Social integration, social networks, social support, and health. In *Social Epidemiology*, ed. L Berkman, I Kawachi, pp. 137–73. New York: Oxford Univ. Press
5. Bertrand JT, O'Reilly K, Denison J, Anhang R, Sweat M. 2006. Systematic review of the effectiveness of mass communication programs to change HIV/AIDS-related behaviors in developing countries. *Health Educ. Res.* 21(4):567–97
6. Borsari B, Carey KB. 2003. Descriptive and injunctive norms in college drinking: a meta-analytic integration. *J. Stud. Alcohol.* 64(3):331–41
7. Boulay M, Storey JD, Sood S. 2002. Indirect exposure to a family planning mass media campaign in Nepal. *J. Health Commun.* 7(5):379–99
8. Brown SS, Eisenberg L, eds. 1995. *The Best Intentions: Unintended Pregnancy and the Well-Being of Children and Families*. Washington, DC: Natl. Acad. Press
9. Cent. Health Commun. 2007. *The Harvard Mentoring Project*. <http://www.hsph.harvard.edu/chc/mentoring.html>
10. Chad-Olmsted SM. 2002. Branding and Internet marketing in the age of digital media. *J. Broadcast. Electron. Media* 46(4):641–45
11. Christensen Clayton M. 1997. *The Innovator's Dilemma: When New Technologies Cause Great Firms to Fail*. Boston: Harvard Bus. Sch. Press
12. Cohen D, Scribner R, Farley T. 2000. A structural model of health behavior: a pragmatic approach to explain and influence health behaviors at the population level. *Prev. Med.* 30:146–54
13. Darnton-Hill I, Nalubola R. 2002. Fortification strategies to meet micronutrient needs: successes and failures. *Proc. Nutr. Soc.* 61:231–41
14. DeJong W, Schneider SK, Towvim LG, Murphy MJ, Doerr EE, et al. 2006. A multisite randomized trial of social norms marketing campaigns to reduce college student drinking. *J. Stud. Alcohol.* 67(6):868–79
15. Dorfman L, Wallack L, Woodruff K. 2005. More than a message: framing public health advocacy to change corporate practices. *Health Educ. Behav.* 32(3):320–36
16. DuRant RH, Wolfson M, LaFrance B, Balkrishnan R, Altman D. 2006. An evaluation of a mass media campaign to encourage parents of adolescents to talk to their children about sex. *J. Adolesc. Health* 38(3):298.e1–9
17. Durantini M, Albarracin D, Mitchell A, Earl A, Gillette J. 2006. Conceptualizing the influence of social agents of behavior change: a meta-analysis of the effectiveness of HIV prevention interventionists for different groups. *Psychol. Bull.* 132:212–48

18. Emmons KM, Barbeau EM, Gutheil C, Stryker JE, Stoddard AM. 2007. Social influences, social context, and health behaviors among working-class, multi-ethnic adults. *Health Educ. Behav.* 34(2):315–34
19. Farley T, Cohen D. 2005. *Prescription for a Healthy Nation: A New Approach to Improving Lives by Fixing our Everyday World*. Boston: Beacon
20. Farrelly MC, Davis KC, Haviland ML, Messeri P, Healton CG. 2005. Evidence of a dose-response relationship between “truth” antismoking ads and youth smoking prevalence. *Am. J. Public Health* 95(3):425–31
21. Farrelly MC, Healton CG, Davis KC, Messeri P, Hersey JC, Haviland ML. 2002. Getting to the truth: evaluating national tobacco countermarketing campaigns. *Am. J. Public Health* 92(6):901–7
22. Fischer CS. 1982. *To Dwell Among Friends*. Chicago: Univ. Chicago Press
23. Fishbein M. 2000. The role of theory in HIV prevention. *AIDS Care* 12:273–78
24. Flora JA, Maibach EW, Maccoby N. 1989. The role of media across four levels of health promotion intervention. *Annu. Rev. Public Health* 10:181–201
25. Flynn BS, Worden JK, Bunn JY, Dorwaldt AL, Dana GS, Callas PW. 2006. Mass media and community interventions to reduce alcohol use by early adolescents. *J. Stud. Alcohol.* 67(1):66–74
26. Glantz SA. 2003. Smoking in movies: a major problem and a real solution. *Lancet* 362(9380):258–59
27. Granfield R. 2002. Believe it or not: examining to the emergence of new drinking norms in college. *J. Alcohol Drug Educ.* 47(2):18–31
28. Grierson T, van Dijk MW, Dozois E, Mascher J. 2006. Using the Internet to build community capacity for healthy public policy. *Health Promot. Pract.* 7(1):13–22
29. Hafstad A, Aarø LE, Engeland A, Andersen A, Langmark F, Stray-Pedersen B. 1997. Provocative appeals in antismoking mass media campaigns targeting adolescents—the accumulated effect of multiple exposures. *Health Educ. Res.* 12(2):227–36
30. Harwood EM, Witson JC, Fan DP, Wagenaar AC. 2005. Media advocacy and underage drinking policies: a study of Louisiana news media from 1994 through 2003. *Health Promot. Pract.* 6(3):246–57
31. Hicks JJ. 2001. The strategy behind Florida’s “truth” campaign. *Tob. Control* 10:1–2
32. Holder HD, Treno AJ. 1997. Media advocacy in community prevention: news as a means to advance policy change. *Addiction* 92(Suppl. 2):S189–99
33. Hornik R. 2002. *Public Health Communication: Evidence for Behavior Change*. Mahwah: Erlbaum
34. House J, Landis K, Umberson D. 1988. Social relationships and health. *Science* 241:540–45
35. Huhman ME, Potter LD, Duke JC, Judkins DR, Heitzler CD, Wong FL. 2007. Evaluation of a national physical activity intervention for children: The VERB Campaign 2002–2004. *Am. J. Prev. Med.* 32(1):38–43
36. Inst. Med. 2002. *Speaking of Health: Assessing Health Communication Strategies for Diverse Populations*. Washington, DC: Natl. Acad. Press
37. Inst. Med. 2003. *Who Will Keep the Public Healthy? Educating Public Health Professionals for the 21st Century*. Washington, DC: Natl. Acad. Press
38. Isen A. 2001. An influence of positive affect on decision making in complex situations: theoretical issues with practical implications. *J. Consum. Psychol.* 11:75–85
39. Kahn EB, Ramsey LT, Brownson RC, Heath GW, Howze EH, et al. 2002. The effectiveness of interventions to increase physical activity: a systematic review. *Am. J. Prev. Med.* 22(S1):73–107

40. Kawachi I, Kennedy BP, Glass R. 1999. Social capital and self-rated health: a contextual analysis. *Am. J. Public Health* 89(8):1187–93
41. Kawachi I, Kennedy BP, Lochner K, Prothrow-Stith D. 1997. Social capital, income inequality, and mortality. *Am. J. Public Health* 87(9):1491–98
42. Lefebvre RC. 2006. *The implications of “social media” for social marketing and public health*. Lecture at George Washington Univ., Sept. 27
43. Maibach E. 2007. The influence of the media environment on physical activity: looking for the big picture. *Am. J. Health Promot.* 21:S353–62
44. Maibach E, Abrams L, Marosits M. 2007. Communication and marketing as tools to cultivate the public’s health: a proposed “people and places” framework. *BMC Public Health* 7:88–102
45. Maibach E, Parrott R. 1995. *Designing Health Messages: Approaches from Communication Theory and Public Health Practice*. Thousand Oaks, CA: Sage
46. Marmot MG, Smith GD, Stansfeld S, Patel C, North F, et al. 1991. Health inequalities among British civil servants: the Whitehall II study. *Lancet* 337:1387–93.
47. McKinney M, Banwart M. 2005. Rocking the youth vote through debate: examining the effects of a citizen versus journalist controlled debate on civic engagement. *Journal. Stud.* 6:153–63
48. McLeroy KR, Bibeau D, Steckler A, Glanz K. 1988. An ecological perspective on health promotion programs. *Health Educ. Q.* 15:351–77
49. Mitchell E. 1997. The changing epidemiology of SIDS following the national risk reduction campaigns. *Pediatr. Pulmonol. Suppl.* 16:117–19
50. Montgomery KC. 1993. The Harvard Alcohol Project: promoting the designated driver on television. In *Organizational Aspects of Health Communication Campaigns: What Works?* ed. T Backer, E Rogers, pp. 178–213. Newbury Park, CA: Sage
51. Niederdeppe J, Farrelly MC, Wenter D. 2007. Media advocacy, tobacco control policy change and teen smoking in Florida. *Tob. Control* 16(1):47–52
52. Noar SM. 2006. A 10-year retrospective of research in health mass media campaigns: Where do we go from here? *J. Health Commun.* 11(1):21–42
53. O’Donnell MP. 2005. A simple framework to describe what works best: improving awareness, enhancing motivation, building skills, and providing opportunity. *Am. J. Health Promot.* 20(1):suppl 1–7 following 84, iii
54. Off. Natl. Drug Control Policy (ONDCP). 2002. *Evaluation of the National Youth Anti-Drug Media Campaign: fifth semi-annual report of findings*. <http://www.mediacampaign.org/publications/westat5/pdf.html> on 5/1/07
55. Palmgreen P, Donohew L, Lorch E, Hoyle R, Stephenson M. 2001. Television campaigns and adolescent marijuana use: tests of sensation seeking targeting. *Am. J. Public Health* 91:292–96
56. Pierce JP, Gilpin EA, Emery SL, White MM, Rosbrook B, et al. 1998. Has the California tobacco control program reduced smoking? *JAMA* 280(10):893–99
57. Putnam RD. 1995. Bowling alone: America’s declining social capital. *J. Democr.* 6(1):65–78
58. Ram R. 2006. Further examination of the cross-country association between income inequality and population health. *Soc. Sci. Med.* 62:779–91
59. Randolph W, Viswanath K. 2004. Lessons learned from public health mass media campaigns: marketing health in a crowded media world. *Annu. Rev. Public Health* 25:419–37
60. Reynolds JH, Hobart RL, Ayala P, Eischen MH. 2005. Clean indoor air in El Paso, Texas: a case study. *Prev. Chron. Dis.* 2(1):1–7

61. Roberts DF, Foehr UG, Rideout MA. 2005. *Generation M: Media in the Lives of 8–18 Year Olds*. Menlo Park, CA: Kaiser Family Found.
62. Rogers E. 2003. *Diffusion of Innovation*. New York: Free Press. 5th ed.
63. Rogers EM, Storey JD. 1987. Communication campaigns. In *Handbook of Communication Science*, ed. C Berger, S Chaffee, pp. 817–46. London: Sage
64. Ross MW, Rigby K, Rosser BR, Anagnostou P, Brown M. 1990. The effect of a national campaign on attitudes toward AIDS. *AIDS Care* 2(4):339–46
65. Russell CA, Clapp JD, Dejong W. 2005. Done 4: analysis of a failed social norms marketing campaign. *Health Commun.* 17(1):57–65
66. Salis J, Cervero R, Ascher W, Henderson K, Kraft K, Kerr J. 2006. An ecological approach to creating active living communities. *Annu. Rev. Public Health* 27:1–14
67. Sargent JD, Beach ML, Adachi-Mejia AM, Gibson JJ, Titus-Ernstoff LT, et al. 2005. Exposure to movie smoking: its relation to smoking initiation among US adolescents. *Pediatrics* 116(5):1183–91
68. Shefner-Rogers CL, Sood S. 2004. Involving husbands in safe motherhood: effects of the SUAMI SIAGA campaign in Indonesia. *J. Health Commun.* 9(3):233–58
69. Siegel M, Biener L. 2000. The impact of an antismoking media campaign on progression to established smoking: results of a longitudinal youth study. *Am. J. Public Health* 90(3):380–86
70. Slater MD. 2006. Specification and misspecification of theoretical foundations and logic models for health communication campaigns. *Health Commun.* 20(2):149–57
71. Sly DF, Hopkins RS, Trapido E, Ray S. 2001. Influence of a counteradvertising media campaign on initiation of smoking: the Florida “truth” campaign. *Am. J. Public Health* 91(2):233–38
72. Snyder LB, Hamilton MA. 2002. A meta-analysis of U.S. health campaign effects on behavior: emphasize enforcement, exposure, and new information, and beware the secular trend. In *Public Health Communication: Evidence for Behavior Change*, ed. RC Hornik, pp. 357–84. Mahwah, NJ: Erlbaum
73. Sparks C. 2007. Advocacy as a tobacco control strategy. In *Ending the Tobacco Problem: A Blueprint for the Nation*, ed. RJ Bonnie, K Stratton, RB Wallace, pp. 690–703. Washington, DC: Natl. Acad. Press
74. Stead M, Hastings G, Eadie D. 2002. The challenge of evaluating complex interventions: a framework for evaluating media advocacy. *Health Educ. Res.* 17(3):351–64
75. Stillman FA, Cronin KA, Evans WD, Ulasevich A. 2001. Can media advocacy influence newspaper coverage of tobacco: measuring the effectiveness of the American stop smoking intervention study’s (ASSIST) media advocacy strategies. *Tob. Control* 10(2):137–44
76. Stokols D. 1992. Establishing and maintaining health environments: toward a social ecology of health promotion. *Am. Psychol.* 47:6–22
77. Strecher V. 2007. Internet methods for delivering behavioral and health-related interventions (eHealth). *Annu. Rev. Clin. Psychol.* 3:53–76
78. Suggs LS. 2006. A 10-year retrospective of research in new technologies for health communication. *J. Health Commun.* 11:61–74
79. Surkan PJ, Dejong W, Herr-Zaya KM, Rodriguez-Howard M, Fay K. 2003. A paid radio advertising campaign to promote parent-child communication about alcohol. *J. Health Commun.* 8(5):489–95
80. Thorson E, Beaudoin CE. 2004. The impact of a health campaign on health social capital. *J. Health Commun.* 9(3):167–94

81. Triplett W. 2007. Anti-smoking campaign underway. *Variety*, Jun. 22. <http://www.variety.com/article/VR1117967461.html?categoryid=1064&cs=1>
82. Umberson D. 1987. Family status and health behaviors: social control as a dimension of social integration. *J. Health Soc. Behav.* 28(3):306–19
83. U.S. Census Bur. 2000. *Census 2000 summary file*. [http://factfinder.census.gov/servlet/DTTable?\\_bm=y&-geo\\_id=01000US&-ds\\_name=DEC\\_2000\\_SF1\\_U&-mt\\_name=DEC\\_2000\\_SF1\\_U\\_PCT012](http://factfinder.census.gov/servlet/DTTable?_bm=y&-geo_id=01000US&-ds_name=DEC_2000_SF1_U&-mt_name=DEC_2000_SF1_U_PCT012) on 10/1/07
84. Valente T, Hoffman B, Ritt-Olson A, Lichtman K, Johnson A. 2003. Effects of a social-network method for group assignment strategies on peer-led tobacco prevention programs in schools. *Am. J. Public Health* 93:1837–43
85. Valente TW, Saba WP. 2001. Campaign exposure and interpersonal communication as factors in contraceptive use in Bolivia. *J. Health Commun.* 6(4):303–22
86. Villani S. 2001. Impact of media on children and adolescents: a 10-year review of the research. *J. Am. Acad. Child. Adolesc. Psychiatry* 40(4):392–401
87. Viswanath K, Randolph Steele W, Finnegan JR Jr. 2006. Social capital and health: civic engagement, community size, and recall of health messages. *Am. J. Public Health* 96(8):1456–61
88. Wallack L. 2003. The role of mass media in creating social capital: a new direction for public health. In *Health and Social Justice: Politics, Ideology, and Inequity in the Distribution of Disease*, ed. R Hofrichter, pp. 594–625. San Francisco: Jossey-Bass
89. Winett LB, Wallack L. 1996. Advancing public health goals through the mass media. *J. Health Commun.* 1(2):173–96
90. Woodruff K. 1995. Media strategies for community health advocacy. *Prim. Care* 22(4):805–15
91. Woodruff K, Dorfman L, Berends V, Agron P. 2003. Coverage of childhood nutrition policies in California newspapers. *J. Public Health Policy* 24(2):150–58
92. Worden JK, Flynn BS. 2000. Effective use of mass media to prevent cigarette smoking. *J. Public Health Manag. Pract.* 6(3):vii–viii
93. Zimmerman FJ, Bell JF. 2006. Income inequality and physical and mental health: testing associations consistent with proposed causal pathways. *J. Epidemiol. Community Health* 60:513–21



**Figure 1**  
A People and places framework for public health.